

Index

Note: Page numbers followed by “f” denotes figure and “t” denotes table respectively.

a

- abdominal cavity 576–579
- abdominal fluid 574
- abdominal viscera, pathology of 563–568
- abnormal navicular bone flexor surface 40f
- abscess 211, 213f, 214, 540
 - aortic root 438f
 - dorsal carpus 137f
 - of heterogenous echodensity 418
 - internal 462f, 463f
 - laryngeal lymph node 522f
 - mesenteric 562, 563f
 - neck 538f, 539f
 - perirectal 480f
 - perivascular 544f
 - pleural 415, 415f
 - pulmonary 418, 418f
 - retrobulbar 487
 - retropharyngeal lymph node 523f, 524f
 - stromal 484, 489f, 490f
 - subsolar 35f
- absorption 3
- accessory carpal bone (ACB) 130, 131, 132f
- accessory ligament of the deep digital flexor tendon (ALDDFT) 89, 90, 96
 - desmitis 118, 120, 120f
 - distal insertional desmopathy 121f
 - fore limb 122f
 - hind limb 120, 122
- accessory ligament of the SDFT (AL-SDFT) 89, 132, 133f
 - desmitis 140f
- acepromazine, use in separating unilateral twins 364
- acetabulum 235, 236, 236f, 237f
- acoustic coupling gel (ACG) 30, 409
- acoustic enhancement artifact 14–15, 14f
- acoustic impedance 2–3, 2t
- acoustic shadowing artifact 15, 15f
- acute hepatitis 449f
- acute synovitis 55, 56f
- acute tendinitis, of biceps brachii tendon 162f, 164f
- adenocarcinoma 439f
- adhesion 76
 - bladder mucosal 460f
 - deep digital flexor tendon 31, 39, 73
 - spleen and stomach 453f
- adjacent medulla 446
- adult:
 - bladder in 458f
 - colitis in 476f, 477f

horse, enteritis in 472f
 aerophagia 569f
 air, in uterus 339, 339f
 air-soft-tissue interface 4
 allantoic cavity 354f
 amniotic fluid 390
 A-mode, equine ocular ultrasonography 482
 ampullae 271
 occluded 297
 per rectum 290
 plugged 286
 of vasa deferentia 276f–280f, 294f, 295f
 of vas deferens 283f–284f
 ampullar lumen 285f
 amyloidosis 449f
 anechoic fluid 419
 annular ligament syndrome (ALS) 76–77
 anovulatory follicles 332–333, 332f, 333f
 antebrachiocarpal joint (ACJ) 128
 anterior chambers 483
 fibrin accumulation in 485
 synechiae 484–485
 anterior lens luxation 499f, 500f
 Anti-Müllerian Hormone (AMH) 333
 aorta 386, 546f
 cranial chest and 386f
 vascular structures 540, 546
 aortic diameter:
 fetus, 234 days gestation 384, 386f
 aortic regurgitation 434f, 435f, 436f, 441f
 aortic root
 abscess 438f
 rupture 443f
 aortic valve closure (AVC) 441f
 aortic valve lesions 433f
 aphakia 485
 arteriovenous fistula 544f
 arthritis 55–57
 arthroscopy 84
 articular cartilage 256
 articular long bone fractures 67
 artifacts 11–17, 488
 absorption technique 488
 acoustic enhancement 14–15, 14f
 acoustic shadowing 15, 15f

B-line 13, 14, 14f
 comet-tail 12, 13, 14, 411f, 412, 413
 definition 11
 edge artifact, blood vessel 230
 edge shadowing 15, 16f
 eye, Baum's Bump 488
 mirror-image 17
 operator error 11–12
 reverberation 12–14, 12f, 13f
 reverberation technique 488, 488f
 ring-down 12, 13, 14f
 side-lobe 16–17, 17f
 slice-thickness 15–16, 16f
 artifactual sediment 16
 asteroid hyalosis 486
 asystole 388
 atlanto-occipital (AO) joint 249, 251f
 assessment 250
 composite transverse image 251f
 longitudinal image 251f, 252f
 synovial sepsis 250
 attenuation 2
 causes of 2
 auriculo-palpebral nerve block 482
 axial resolution 5, 6f

b

back
 caudal thoracic and lumbar facet
 joints 245–248
 epaxial musculature 248
 indications 243
 interspinous ligaments 243–245
 supraspinous ligaments 243–245, 244f, 245f
 ultrasound assessment 243
 bacterial peritonitis 460f, 461f, 462f
 bandage bows 103, 105, 105f
 basihyoid bone, lingual process of 258
 Bedside Ocular Ultrasound 481
 biceps brachii tendon 149, 161–163
 acute tendinitis 162f, 164f
 distal insertion 155, 158f
 enthesopathy 155
 normal proximal 161f
 origin 160f

proximal 164f
 bicipital bursa 155, 162, 163, 166
 bicipital bursitis 161
 chronic 165f, 166f
 proximal humeral osteitis and 166f
 septic 163
 bicipital tendinitis 161, 163
 bile duct stricture 450f, 451f
 biophysical profile score (BPS) 403
 bladder 455–458
 adult 458f
 with cystitis 564f
 foal 458f
 normal 579f
 rupture 459f, 562–563, 564f
 with sediment 459f
 urinary calculi 460f
 urine-filled 562
 uroperitoneum 460f
 wall defect 459f
 “B-line” artifacts 13, 14, 14f
 blood (ifema) 485
 blunt force trauma 560
 blunt trauma 505f
 B-mode 3–4
 equine ocular ultrasonography 482–483
 bone cyst. *See* cyst-like lesions
 bones 134–136, 143, 146
 fragment 63f
 neck 249–254
 bone-soft-tissue interface 3, 4, 4f
 bony injuries 122–126
 bony structures
 tarsus 181, 183
 brachialis muscle 149
 brachiocephalic muscle forms 537
 brain sparing effect 403
 broad ligament hematoma 563, 565f
 bronchopneumonia 574f
 broodmare, pregnant 545f
 “bucket-handle” tear 217f, 218
 bulbospongiosus muscle 300
 bulbourethral glands 271, 272, 285
 color Doppler ultrasound image 294f

 in sexually rested stallion 293f
 ultrasound image of 293f
 bursae 143
 buscopan (N-butylscopolammonium bromide) 323, 347, 365, 372
C
 calcaneal bursa 180
 calcaneal tendon 174
 calcification 77
 biceps brachii tendon 162
 calcinosis circumscripta 226
 capsular rupture 225
 cardiology. *See also* echocardiography
 normal equine heart, ultrasonographic anatomy of 422, 426
 technical considerations 421–422
 carotid artery 518
 jugular groove 532
 jugular vein and 529f–530f, 532
 carpal canal (CC) 131, 132f, 133f, 143f
 carpometacarpal joint (CMCJ) 128
 carpus
 anatomy and scanning technique 128–134
 bones 143, 146
 dorsal 128–129, 129f
 joints and bone 134–136
 joint surfaces 146
 lateral 129–130, 130f, 131f
 medial 130–131, 131f, 133f
 palmar 131–134, 134f, 135f
 superficial soft tissues 143
 surgery 146
 tendons and ligaments 143
 ultrasonographic abnormalities 136–146
 cartilage dissection 60
 cataracts 486
 congenital 496f
 and uveitis 497f
 caudal cervical vertebrae 253
 caudal cruciate ligament (CaX) 198f, 200, 201f, 203
 caudal horn tears 218, 219f
 caudal thoracic joints 245–248
 caudal vaginal 560

- cecal impaction 478, 479f
- cecal inversion 478, 479f
- cecocolic intussusception 478, 479f
- cecum 465, 478, 559, 570, 578f
- cellulitis
 - focal 540
 - perivascular 540f
- central process 300
- cephalic vein (CV) 132
- cervix 325, 342
 - abnormal 329f
 - after ovulation 328f
 - diestrus 329f
 - during early estrus 326f
 - in early pregnancy 348
 - estrus, failure to relax in 342
 - intrauterine fluid adjacent to 343f
 - ovulation approaches 327f
 - 250 days gestation 392f
 - 244 days gestation 393f
 - 277 days gestation 393f
 - 232 days gestation 392f
 - vagina and 345f
 - ventral 344f
- chip fracture
 - distal craniomedial radius 145f
 - multiple palmar 146f
- chip fractures 61, 63
- choke 534
- cholangiohepatitis 450f, 451f
- cholelithiasis 451f
- chondromalacia patellae 210
- chorioallantoic membranes 357
- choroid 484
- chronic acetabular fracture 238f
- chronic collateral ligament desmitis 66
- chronic hypertrophic synovitis 57, 58f, 59
- chronic laminitis 37f
- chronic lymphangitis 544f
- chronic scapular fracture 159f
- chronic synovitis 56, 203, 210
- chronic tendinopathy 100–102, 101f
- ciliary body cysts 485
- “cine recall system” 386
- circumflexa ilium profunda artery 400
- clitoris, in female 370
- cold effusion 55, 203
- colitis 578f
 - in adult 476f, 477f
 - Potomac Horse Fever 477f
 - right dorsal 475f, 476f
 - from Salmonellosis 476f
- collateral ligaments (CL):
 - complete rupture 64
 - desmitis 174
 - desmitis and tears 215
 - desmopathy 64
 - diffuse 65f
 - of distal interphalangeal joint 26, 28f
 - of elbow joint 149–150
 - hock 173–174
 - injury in horse 65f
 - medial 64, 65f
 - structure 54f
- collateral sesamoidean ligament. *See*
 - “T” ligament
- colliculus masculinus:
 - cross-sections of 295f
 - cysts of 286
 - utriculus masculinus in 286
- colliculus seminalis 272, 273
 - midline cyst of 275f–276f
- colon:
 - gas-distended 473, 474f
 - large 569
 - left ventral 474, 474f
 - lumen of 476f
 - right dorsal 473, 474f
 - right dorsal displacement of 475, 475f
 - small 570
 - torsion 477f, 478f
 - torsion or volvulus of 476f
 - ventral 559
 - wall 475f
- color Doppler 482
 - in horses 426f, 433f
 - portal vein 448f
- color flow Doppler 19, 20f, 421, 429
 - of luminal blood flow 543f
- colt. *see* fetus

- combined thickness of the utero-placental unit (CTUP) 384, 393–394
 - comet-tail artifact 12, 13, 14, 411f, 412, 413
 - common arterial trunk (CAT) 431f
 - common carotid artery, bifurcation of 520f
 - common digital extensor (CDE) tendon 52, 60f
 - common digital extensor tendon (CDET) 90, 129, 138f, 142f
 - congenital cataracts 496f
 - console 5
 - continuous-wave (CW) Doppler 19–20, 19f, 421, 429, 430f, 433f, 437f
 - contralateral ovary 333, 334
 - contrast tenography 43
 - convex probes 7
 - core lesions 205
 - cornea 483
 - corneal anesthesia 482
 - corneal damage 482
 - corneal edema 484, 489f, 492f
 - corneal ulcer 484, 494f
 - corpa lutea 349f
 - corpora cavernosa 300, 302
 - corpora lutea (CLs) 348
 - corpora nigra* 483, 485
 - cor pulmonale 442f
 - corpus cavernosum 302, 305f
 - dorsal aspect 305f
 - processes 306f
 - trabeculae of 304f, 308f, 309f
 - corpus hemorrhagicum 328f, 334
 - corpus luteum 326, 329f, 330f, 348, 349f
 - corpus spongiosum 302
 - glandis 300
 - trabeculae in 308f, 309f
 - coxofemoral joint 236f
 - cranial cruciate ligament (CrX) 200, 201f, 205f, 222f
 - complete rupture of 223f
 - “cranial drawer” 220
 - cranial meniscotibial ligaments (CMTL) 202, 215, 220f
 - of medial meniscus 220f
 - tears 218
 - cranial thorax 409
 - cranial tibial muscle 177
 - cranial tibial tendon 177
 - cranioventral abdomen 448f
 - crena marginis solearis 34f
 - cricoarytenoideus lateralis muscle (CAL) 259, 260f
 - cross-sectional area (CSA) 91, 93, 96
 - cruciate ligament (XL) injuries 220
 - cryptorchid 286f, 299
 - crystalline 483
 - crystalline lens 485, 486
 - dislocation 486
 - crystalline material 451f
 - cunean bursa 180
 - cunean tendon (CT) 177, 182f
 - curvilinear transducers 7
 - cystitis 562
 - bladder with 564f
 - sabulous 564f
 - with urinary calculi 564f
 - cyst-like lesions 59–61
 - cysts
 - ciliary body 485
 - endometrial 335, 337f, 338f, 339, 361–362
 - epithelial inclusion 334, 334f
 - iris 485, 506f
 - lymphatic 335
 - parovarian 334, 334f
 - uterine 350
- d**
- daily sperm output (DSO) 317, 318
 - deep digital flexor tendon (DDFT) 25, 29, 31, 37f, 38f, 39f, 69, 72, 73, 73f, 77f, 86, 89, 90
 - accessory ligament of 123f
 - chronic 78f
 - cross-sectional areas 91
 - lateral 175, 180
 - lesions 35
 - marginal tears detection 78f, 79f
 - medial 175, 180, 183f
 - mid-substance tendinopathy 77f
 - over-strain midsubstance injuries 77
 - septic carpal canal and 143f

- tears 79–80
- tendinopathy/desmopathy 106
- tendonitis of 177
- transverse image of 182f
- transverse transverse ultrasound of 33f
- trauma 80
- deep distal sesamoidean ligaments (DDSL) 67
- depth 8
- descemetocele 492f
- desmitis 92, 106, 144f
 - accessory ligament of the DDFT 118–120, 120f
 - accessory ligament of the SDFT 140f
 - body of SL 113–117, 115f, 116f
 - collateral ligaments 174, 215
 - of long plantar ligament 108f, 175
 - of medial PL 205
 - proximal suspensory 107–113, 109f, 110f, 111f
 - suspensory ligament branches 117–118
- detomidine 323
- diagnostic ultrasonography 5
- diaphragmatic hernia 559–560, 560f
- diastolic left ventricular function 429
- diestrus 348
 - cervix 329f
 - non-pregnant mare in 351
 - uterus 329f, 335
- diffuse collateral ligament injury 65
- diffuse tendinopathy 93, 94
- digital annular ligaments 44, 47
- digital extensor tendons 52
- digital flexor tendinopathy
 - lateral 125f
- digital flexor tendon sheath 68–73, 74f, 75f
 - deep digital flexor tendon
 - abnormalities 77–80, 77f
 - infected, characteristics of 73f
 - manica flexoria 70, 71f, 72
 - normal ultrasonographic anatomy 69–73
 - palmar/plantar annular ligament syndrome 76–77
 - percutaneous trauma to 73f
 - primary tenosynovitis 73–76
 - proximal pouch 69, 70f
 - sepsis of 73
 - superficial digital flexor tendon
 - abnormalities 80–82
- digital sheath tenosynovitis 76
- dissecting lesion 206
- distal interphalangeal (DIP) joint 25, 86
 - abnormalities 29
 - collateral ligaments 26, 28f
 - normal ultrasonographic appearance 28f
 - osteoarthritis 30f
 - palmar/plantar pouch 26
 - ruptured collateral ligament 30f
 - ultrasound-guided injection 28f
- distal interphalangeal joint (DIPJ) 31
- distal limb vasculature 540
- distal radial artery (DRA) 132
- distal sesamoid bone (DSB) 29, 30, 31f
 - flexor surface and distal aspect 31
 - fractures 35
 - hyperechoic flexor cortical surface of 32f
 - proximal displacement 38f
 - severe fibrocartilage degeneration of 39f
- distal sesamoidean impar ligament (DSBIL)
 - fibers 31, 35, 38f
 - hypoechoic appearance 38f
- distal sesamoidean ligaments (DSLs) 41–42, 67
 - abnormalities 42–44
- Doppler. *See also* Doppler imaging
 - carotid artery 403f
 - color 426f
 - color flow 19, 20f, 421
 - continuous-wave 19–20, 19f, 421, 429, 430f
 - echocardiography 429, 433
 - Middle Cerebral Artery 402
 - modes 18
 - power 20
 - pulsed-wave 18–19, 19f, 421, 429
 - right uterine artery 401f
 - scans of maternal uterine arteries 399
 - spectral 18, 18f
 - technology 383
 - umbilical artery 401f, 402f
 - velocimetry parameters 399–400
- Doppler imaging 17–20

- color flow Doppler 19, 20f
- continuous wave Doppler 19–20, 19f
- negative frequency shift 17–18, 18f
- positive frequency shift 17, 18f
- power Doppler 20
- pulsed wave Doppler 18–19, 19f
- spectral Doppler 18, 18f
- Doppler shift 18, 18f, 20
- dorsal carpus 128–129, 129f
 - abscess 137f
 - normal neonate 136f
 - premature foal 136f
- dorsal sacroiliac ligaments 231, 232, 234f, 235, 235f
 - central hypoechogenic regions with 235f
 - normal appearance of 234f, 235f
 - ventral parts of 235
- dorsal spinous processes (DSPs) 244, 245f
- duodenal distension 466, 467f
- duodenal stricture 467f
- duodenitis 466, 467f, 570, 577f
- duodenum 448f, 466, 467f, 468f, 479f, 570
- dystrophic mineralization, neck 249f
- e**
- early pregnancy examination 358
- echocardiogram 436f
- echocardiography 426. *See also* cardiology
 - detection 428
 - Doppler 429, 433
 - endocardial cushion defect 430f
 - great vessels 424f
 - in horses. *See* horses, echocardiography in
 - left atrial size and function 424f, 428
 - left ventricular size and function 423f, 428
 - methods 429
 - mitral and aortic valve motion 425f
 - restrictive ventricular septal defect 427f
- echogenicity 3–4, 99f
 - of intrauterine fluid 335
- edge refraction. *See* edge shadowing artifact
- edge shadowing artifact 15, 16f
- elbow 148–149
 - abnormalities 148–155
 - biceps brachii muscle 155
 - biceps brachii tendon 155, 161–163
 - collateral ligaments of 149–150
 - craniomedial aspect of 158f
 - dorsal aspect of 149f
 - infraspinatus muscle 166–167
 - lateral collateral ligament of 149–150, 152f, 153f, 154f
 - medial collateral ligament 153f
 - osteoarthritis 149, 151f
 - preparation and scanning technique 148
 - scapulohumeral joint 167–172
 - septic arthritis of 150f
 - supraspinatus muscle and tendons 163–166
 - triceps brachii muscle 155
 - ulnaris lateralis muscle 150, 155, 156f, 157f
 - ultrasonographic anatomy 148–155
- electronic transducers 6
- embryo. *See also* pregnancy
 - day 18 353f, 360
 - day 15 352f, 360
 - day 14 352f, 360, 361f
 - day 17 352f
 - day 13 360f
 - day 32 355f
 - day 20 353f, 354f
 - day 25 355f
 - day 29 355f
 - day 21 354f, 355f, 360f
 - day 22 355f
 - distinguishing from endometrial
 - cysts 361–362
 - pregnancy stage 349
 - embryo proper (EP) 354f, 355f
 - empty trophoblastic vesicle (ETV) 358f
 - endocardial cushion defect 430f
 - endocarditis 426
 - endometrial culture 335
 - endometrial cups 339, 341f, 354
 - endometrial cysts 335, 337f, 338f, 339, 361–362
 - major consequence of 335
 - pregnancy adjacent to 337f
 - ultrasonographic appearance of 337f, 338f
 - endometrial edema 335, 336f, 566f
 - abnormal 360f
 - endometrial glands 335

- endometritis 340*f*
- endophthalmitis 505*f*
- enteritis 577*f*
 - in adult horse 472*f*
 - anterior 472*f*
 - post-partum mare secondary to 571*f*
 - proliferative 471
 - small intestinal volvulus and 577*f*
- enthesophytes 59, 64, 110, 118
- enthesophytosis 44
- envelope 18
- epaxial musculature 248
- epicondyles 149
- epididymal tail 317*f*, 318*f*
- epididymis 310, 312*f*
- epididymitis 313, 315*f*
- epithelial inclusion cysts 334, 334*f*
- E-point-to-septal separation (EPSS) 441*f*
- equine chorionic gonadotropin (eCG)
 - 354, 356*f*
- equine multinodular pulmonary fibrosis 417*f*
- equine ocular ultrasonography (EOUS) 481
 - in A-mode 482
 - anterior and posterior chambers 483
 - anterior chamber synechiae 484–485
 - aphakia 485
 - blood 485
 - in B-mode 482–483
 - cataract 486
 - ciliary body cysts 485
 - cornea 483
 - corneal edema 484, 489*f*
 - corneal infiltrates 484
 - corneal ulcer 484
 - crystalline 483
 - crystalline lens dislocation/
 - sub-luxation 486
 - eyelid edema 487
 - fibrin accumulation 485
 - glaucoma 485, 504*f*
 - intra-ocular mass 487
 - iris and ciliary bodies 483
 - iris cysts 485
 - iris prolapse 485
 - keratitis 484, 489*f*
 - lens rupture 486
 - microphakia 486, 497*f*
 - pus (*ipopion*) 485
 - retina/choroid/sclera 484
 - retina detachment 487, 502*f*, 503*f*
 - retrobulbar area 484
 - retrobulbar diseases 487
 - small eyeball 485
 - stromal abscess 484, 489*f*, 490*f*
 - technique 481–483
 - uveitis 485
 - vitreous chamber 484
 - vitreous opacity 486–487
- equine pregnancy
 - biophysical profile score 403
 - doppler velocimetry parameters of 399–400
 - imaging of 384
 - transabdominal evaluation of 385
- equine recurrent uveitis (ERU) 502*f*,
 - 503*f*, 504*f*
- equine reproductive tract 331
- equine shoulder 155
- equipment, pelvis 229
- erect penis, ultrasound evaluation of 302,
 - 303*f*
- esophagus 465–466
 - choke 535*f*
 - fistula 537*f*
 - with food bolus 535*f*
 - jugular groove 534, 537
 - with nasogastric tube 535*f*
 - normal 466*f*, 534*f*
- estrous edema 324*f*, 327*f*
- estrus, failure to relax in 342
- excessive pressure 11
- exophthalmos 508*f*
- extensive hematoma 561*f*
- extensor carpi radialis bursa (ECRB) 129
- extensor carpi radialis tendons (ECRT) 129
- extensor tendons 67, 90, 122
- external carotid 388
- external carotid artery 518
- external genitalia 371, 379
 - after 150 days of Gestation 371
 - 90-150 days 377

- post 150+ days 380
- on transabdominal scan 371
- external iliac artery 563
- external jugular vein 540
- external trauma 534
- external umbilical remnant 547, 549f
- exudative peritonitis 462f
- eye
 - normal 488f
 - volume, fetal 385–386
- eyelid edema 487
- f**
 - facies flexoria 31
 - fascicular pattern 86
 - feed material 534
 - female
 - clitoris in 370
 - genital tubercle for 373
 - femoropatellar joint 191–194
 - abscesses 211, 214
 - degenerative joint disease 208f
 - effusion 203–204
 - hematomas 211, 214
 - inflammatory changes in 207f
 - non-inflammatory effusion of 206
 - osteocondrosis 206, 209
 - patellar fractures 209–210
 - patellar ligament injuries 204–206, 207f
 - stifle 191–194, 191f
 - femorotibial joints 214–226
 - caudal aspect 202–203
 - collateral ligament desmitis and tears 215
 - cranial aspect of 199–202, 199f
 - cruciate ligament injuries 220
 - lateral 195, 198f
 - long digital extensor muscle 225
 - medial 194–195, 197f, 224f
 - medial aspect 216f
 - meniscal injuries 215, 218, 220
 - miscellaneous 225–226
 - osseous cyst-like lesions 220
 - osteoarthritis 220–221
 - peroneus tertius tendon 225
 - septic arthritis 221, 225
 - fetal activity 387, 388–389

- hydroallantois and 391
- fetal cardiac rhythm 388
- fetal development and monitoring. *See also*
 - equine pregnancy; fetus
 - activity 387, 388–389
 - adequate environment 390
 - adequate growth and development 385
 - aortic diameter 386, 386f
 - biophysical profile score 403
 - biparietal diameter 386, 387f
 - combined thickness of the utero-placental unit 393–394
 - equine gravid cervix, assessment of 391–393
 - equipment 384
 - eye volume 385–386
 - fetal circulation, assessment of 401
 - fetal heart rate 387–388
 - FHR reactivity 387–388
 - intracranial circulation 402–403
 - maternal circulation, assessment of 400–401
 - orbital diameters 385–386
 - other organs 394–396
 - peripheral pulses 388
 - pregnancy database 383–384
 - responsive patterns 387
 - transabdominal approach 385
 - transrectal approach 384
 - umbilical circulation 401
 - umbilical cord 396–399
- fetal eye 385–386
 - measurements 385, 386f
- fetal heart rate (FHR) 387–388
 - accelerations 388
- fetal hypoxia 403
- fetal movements 357
- fetal orientation 390
- fetal sex determination 368. *See also* fetus,
 - colt; fetus, filly
 - 55–90 days 370–371
 - 90–150 days 371
 - plane I 372f
 - plane II female 372f
 - plane III 373f

- plane II male 372f
- procedure at 55–90 days 372–377
- procedure 90–150 days 377–379
- stages of gestation 370–372
- take home hints 372
- transabdominal scan post 150+ days,
 - procedure for 379–382
- ultrasonographic planes 372
- fetal sexing. *See* fetal sex determination
- fetal size, aortic diameter 385
- fetal well-being 383
 - assessment 385
 - and optimal development 383
 - sensitive indicators of 387
 - transrectal field evaluation 387
- fetlock 86
 - acute synovitis 55, 56f
 - articular long bone fractures 67
 - bone fragment 63f
 - chronic hypertrophic synovitis 57, 58f
 - chronic synovitis 56
 - cyst-like lesions 59–61
 - deep digital flexor tendon
 - abnormalities 77–80, 77f
 - digital flexor tendon sheath. *See* digital flexor tendon sheath
 - dorsal aspect of 52, 58f, 70f
 - dorso-medioproximal aspect of 64
 - four quadrants of 50f
 - hemarthrosis 56f
 - intersesamoidean ligament. *See* intersesamoidean ligament
 - normal anatomy 51–55
 - osteoarthritis 59
 - osteocondral fragments 61–64
 - osteocondrosis 59–61, 61f, 62f
 - palmar aspect 53
 - palmar/plantar annular ligament of
 - 72, 72f
 - palmar/plantar P1 fragments 67
 - periarticular injuries 67
 - plantar aspect of 51
 - plantaro-abaxial aspect 68f
 - preparation and scanning technique 49–51
 - proximal scutum abnormalities 82–84
 - proximal sesamoid bones fractures 65,
 - 66–67
 - schematic organization of 51f
 - septic arthritis 67–68
 - subchondral bone injury 57–59
 - superficial digital flexor tendon
 - abnormalities 80–82
 - synovitis/arthritis 55–57
 - traumatic arthritis 57–59
- fetus. *See also* fetus, colt; fetus, filly
 - in anterior presentation 390
 - bi-parietal diameter, 121 days
 - gestation 387f
 - day 50 357f
 - day 51 357f
 - day 48 357f
 - day 60 358f
 - neck 390
 - 180-days-old 395f
 - 158-days-old 395f
 - 152-days-old 390
 - 110-days-old 395f
 - 121 days gestation 387f
 - 127-days-old 396f
 - orbital diameter, 211 days gestation 387f
 - 218-days-old 395f
 - 211 days gestation 386f
 - 215-days-old 386f
 - 214-days-old 395f
 - 244 days old 387f, 397f
 - 243-days-old 390
 - 279-days-old 388f
 - 268 days 391f
 - 268-days-old 387f
 - 234-days-old 386f
 - 236-days-old 389f
 - 224-days-old 388f
 - umbilical vein 550
- fetus, colt
 - axial transverse plane of 374f
 - external genitalia. *See* external genitalia
 - female, mammary gland and teats of 371
 - genitalia 370
 - homogenous texture with central vein
 - 380f, 381f

- 95 days of gestation 377f
 - 94 days of gestation 377f
 - 115 days of gestation 376f
 - 109 days of gestation 379f
 - 117 days of gestation 377f
 - 113 days of gestation 375f
 - 130 days of gestation 380f
 - 65 days of gestation 374f
 - fetus, filly 376f, 378f
 - 135-days-old 396f
 - 124-days-old 396f
 - 105 days of gestation 378f
 - 110 days of gestation 378f
 - 134 days of gestation 379f
 - 124 days of gestation 379f
 - 129 days of gestation 379f
 - 122 days of gestation 378f
 - 78 days of gestation 375f
 - 63 days of gestation 375f
 - FHR reactivity 387–388
 - fibrin 414, 495f
 - accumulation 485
 - adhesions 453f
 - peritonitis 459, 461f
 - pleuritis 413f
 - fibularis tertius 225
 - fluid effusion 214
 - foals
 - bladder in 458f
 - chest for thoracic ultrasonography 574
 - haircoat 573
 - heart 573
 - hock 186
 - liver 452f
 - recumbent 573
 - umbilical hernia in 556f
 - focal cellulitis 540
 - focal extensor tendinitis 69f
 - focus 8–9, 9f
 - foot 25
 - distal interphalangeal joint. *See* distal interphalangeal (DIP) joint
 - dorsolateral/dorsomedial aspects 27f
 - palmar aspect 26f
 - proximal to coronary band 25–29
 - transsolar and transcuneal ultrasonography 29–39
 - ultrasonographic abnormalities 26, 29
 - fracture(s)
 - chronic acetabular 238f
 - detection 229
 - development, stages of 231
 - “green stick” 230
 - pelvis, common sites detected *per rectum* 238f
 - rib 412, 412f, 413f
 - scapula body 155, 157
 - stress 229, 243
 - tuber coxa* 233f
 - frame rate 9
 - free fluid 453f
 - frequency 482
 - ultrasound waves of 5, 8
 - fundamental frequencies 9
 - fungal plaque forms 514
- g**
- gain control 8
 - gas decomposition 568f
 - gas echoes 562
 - gastric distention 467f
 - gastric rupture 461f
 - gastrocnemius tendon (GCN) 174
 - superficial digital flexor tendon and 181f
 - gastrointestinal tract 13, 568–571
 - cecum 570
 - duodenal distension 467f
 - duodenitis 467f, 570
 - duodenum 466, 467f, 468f
 - esophagus 465–466, 466f
 - jejunum 570, 570f
 - large colon 569
 - large intestine 473–480
 - mesenteric rent 571
 - rectum 480
 - small colon 570
 - small intestine 466–473, 469f
 - stomach 466, 467f, 570, 571f
 - gelding 292f
 - genioglossus muscles 261
 - genitalia 370, 372

genital tubercle 370, 371
 for male and female 373
 gestation. *See also* fetus; fetus, colt; fetus, filly
 after 150 days 371–372
 80 days of 371, 371f
 55–90 days of 370–371
 90–150 days of 371
 121 days 387f
 100 days of 371f
 211 days 386f
 277 days 391f
 212 days 387f
 gestational age 347
 eye volume and 385
 resistive index for 397f
 gestation, first 60 days of. *See also* pregnancy
 early embryonic imaging 349–358
 endometrial cysts 361–362
 transrectal technique 347–349
 glandular endometrial cysts 361
 glandular parenchyma 516f
 glans penis 300, 305f, 306f
 glaucoma 485, 500f, 504f
 granulomatous lung disease 417
 granulosa cell tumor (GCL) 333–334, 333f
 gravid uterus 384
 “green stick” fractures 230
 “ground glass” appearance 419
 guttural pouches 514
 empyema 518f, 519f
 lateral compartment of 516f
 medial compartment of 516f, 517f
 tympany 517f

h

harmonic waves 9
 head:
 assessment of structures 262–263, 263f
 larynx 257–260, 258f, 259f
 salivary glands 262
 temporomandibular joint 255–257, 256f, 257f
 tongue 260–262, 261f, 262f
 ultrasonographic assessment 255
 ultrasound guided procedures 263–264
 healing, assessment of 97–100

heart 409
 by echocardiography 409
 hemangiosarcoma 420, 453f
 hemarthrosis 56, 56f, 209
 hematomas 209, 211, 214, 561f, 563f
 broad ligament 563, 565f
 mural 563, 566f
 hemopericardium 415f
 hemoperitoneum 463f, 561
 hematoma formation during resolving 561f
 hemorrhage 203, 208f, 339, 415, 486, 564
 periparturient 561
 perirenal 560
 source of 564
 hemothorax 414f, 415
 pleural abscess 415, 415f
 pyothorax 415, 415f
 hepatitis 449f
 hepatoblastoma 452f
 herniation 419
 high suspensory disease. *See* proximal
 suspensory desmitis
 hip joint 235–238, 236f
 osteoarthritis 237f
 hippomane 391, 397f
 hock:
 collateral ligaments 173–174
 foals 186
 intra-arterial perfusion 185–186
 lateral digital extensor tendon 179f
 subcutaneous bursa 180
 subcutis 185
 tendons and ligaments 173–177
 homogenous tissues 3
 horizontal tears 218
 horses, echocardiography in 426
 aortic root rupture 443f
 aortic valve lesions 433f
 apical muscular ventricular septal
 defect 428f
 chamber dimensions 428
 color Doppler in 426f
 common arterial trunk 431f
 cor pulmonale 442f
 diastolic left ventricular function 429

- dilated cardiomyopathy 441f
- hemodynamic assessment 429
- infective endocarditis 438f
- left atrial function 429
- mitral regurgitation 431f–433f
- myocarditis 439f
- non-restrictive ventricular septal defect 427f
- papillary muscle ischemia 442f
- pericardial effusion 438f
- restrictive ventricular septal defect 427f
- systolic left ventricular function 428–429
- valvular regurgitation 433–444
- humeral tubercles 157–161
- hydronephrosis 457f
- hydroureteronephrosis 457f
- hyperechogenic fluid 335
- hyperechogenic foci 415
- hyperechogenic fragments 61
- hyperechoic concretions 286
- hyperechoic shadow 413
- hyperechoic smegma 306f
- hyperplasia 57
- hypervascularity 100
- hyphema 495f
- hypoechoogenicity 97, 110
- hypoechoic muscle 89
- hypoechoic brachiocephalicus muscle 166
- i**
- iliac artery 545f
 - external 563
 - plaque in 546f
 - thrombosis of 540, 546f
 - vascular structures 540, 546
- ilium:
 - pelvis 229–230
 - ultrasonographic assessment of 230f
- image resolution 5–6, 6f
- immature callus 230
- infective endocarditis 438f
- inflammatory changes (synovitis) 203
- infraspinatus bursa 168f
- infraspinatus bursitis 167
- infraspinatus muscle 166–167, 167f, 168f
- injuries:
 - cruciate ligament 220
 - meniscal 215, 218, 220
 - patellar ligament 204–206, 207f
- interarticular ligaments 134
- intercostal spaces (ICS) 409, 410f, 446, 558
- internal abscess 462f, 463f
- internal carotid artery 518
- internal genitalia 272f
 - stallion 285
 - transrectal ultrasound evaluation 272
- internal jugular vein 540
- internal pudendal artery 300
- internal reproductive tract
 - anatomy 271–272
 - normal ultrasonography 272–286
 - palpation *per rectum* 272
- internal umbilical remnant 555f
- internal vaginal rings 272
- intersesamoidean ligament 72
 - abnormalities 82–84
 - desmitis of 65–66
- interspinous ligaments 243–245
- interstitial pneumonia 411f, 419
- intertubercular groove. *See* intertubercular sulcus
- intertubercular sulcus 157–161
- interventricular septum (IVS) 427f
- intestinal tract 558
- intra-arterial perfusion 185–186
- intracranial circulation 402–403
- intraluminal fluid, septic 566f
- intraluminal thrombus 542f
- intra-ocular mass, EOUS 487
- intrauterine fluid 335, 343f, 353f
 - echogenicity of 335
 - uterus 564
- intrauterine growth restriction (IUGR) 385, 402
- intrauterine marbles 339
- intravenous catheter 541f
- inverted uterine horn 566, 567f
- iris cysts 485, 506f
- iris prolapse 485
- ischiocavernous muscles 300

J

Jejunojejunal intussusception 471, 473f

jejunum 570, 571

joint 134–136

capsule 203

coxofemoral 236f

disease 220

hip 235–238, 236f

neck 249–254

surfaces 146

jugular groove 525, 532

carotid artery 529f–530f, 532

esophagus 534, 534f, 535f, 537

trachea 532, 533f, 534, 534f

ultrasound structures in 529f

jugular vein 466f, 529f–533f, 540

internal and external 540

lateral thoracic vein 545f

myositis 541f

normal 541f

occlusion 541f

perivascular cellulitis/edema 544f

thrombophlebitis 542f

thrombosis 542f, 543f

vascular structures 540

juvenile tendinitis 93, 94f

k

keratitis 484, 489f, 499f

kidneys 446, 455

normal left 454f

normal right 454f

Klebsiella pneumoniae 455f

l

large intestine 473–480

larynx 257–260, 258f

anatomy and scanning technique 257–260

lateral transverse view of 259f

ultrasonographic abnormalities 260

ventral aspect of 259f

lateral carpus 129–130, 130f, 131f

premature foal 137f

lateral collateral ligament (LCL) 64, 195

of elbow joint 149–150, 152f, 153f, 154f

hock 173

lateral deep digital flexor tendon 175, 177, 180

lateral digital extensor tendon (LDET) 90, 129, 130f, 179f, 186f

lateral digital flexor tendon (LDFT) 91

lateral femoral condyle (LFC) 198f, 200f

lateral femoral epicondyle (LFE) 198f

lateral femorotibial joint (LFT) 195, 198f, 199f

lateral patellar ligament (LPL) 193f, 206f, 207f, 211f, 212f

lateral region 537, 539

lateral resolution 5, 6f

lateral thoracic vein 545f

lateral tibial condyle (LTC) 198f

lateral trochlear ridge (LTR) 206f, 208f
distal aspect 212f

Lawsonia intracellularis infection 471, 473f

left hemithorax 410

left kidney 446, 558

normal 454f

left testis:

cranial–caudal view of 311f, 313f

lateral view of 311f, 312f

left ventral colon (LVC) 448f, 474f

left ventricular outflow tract (LVOT) 424f

leiomyomas 343f

lens rupture 486, 501f

leptosporosis 456f

lesions 57, 77, 93

meniscal 218

osseous cyst-like 220

ligaments 86–91, 88f

carpus 143

dorsal sacroiliac 232, 234f, 235, 235f

hock 173–177

linear array transducers 384

linear transducers 6, 6f, 27f, 173

standoff for 8f

lipoma 470f, 471f

liver 446, 558, 559

amyloidosis 449f

bile duct stricture 450f

cranioventral abdomen 448f

and duodenum 448f

hepatitis 449f

- normal 447f
- portal vein 448f
- local anesthesia 264, 481
- local traumatic injuries 103–106
- long digital extensor muscle 225
- long digital extensor tendinopathy 124f
- longitudinal tears 218, 219f
- long lateral collateral ligament (LLCL) 173, 174, 176f, 179f
- long medial collateral ligament (LMCL) 174f
- long plantar ligament (LPL) 174, 176f
 - desmitis of 175
- lumbar facet joints 245–248
- lumbar intertransverse joints 240
 - per rectum*, normal 241f
- lumbar intervertebral articulations (IVAs) 245f, 246, 246f, 247f
- lumbosacral joints 238–240
 - per rectum* ultrasonograph 239f, 240f
- lumen 534
- lung:
 - abnormal 411f
 - cardiac notch of 409
 - normal 410f, 411f
 - parenchyma 416f, 417, 419
 - thoracic ultrasonography 410f
- LV peripheral wall (LVPW) 441f
- lymphatic cysts 335
- lymph nodes 520f
- laryngeal 522f
- retropharyngeal 524f
- lymphosarcoma 420, 453f

m

- magnetic resonance imaging (MRI) 25
- male:
 - genital tubercle for 373
 - penis in 370
 - umbilical cord in 370
- mammary gland 377
- mandibular salivary gland 518, 519f
- manica flexoria 70, 71f, 72, 74, 82, 82f
 - partial/complete tearing 80
 - tearing 81f, 83f
- marbles, uterus 339, 342f
- mare reproductive abnormalities:

- cervix 342
- ovaries. *See* ovary/ovaries
- ultrasound examination 331
- uterus 335–342
- vagina 342–345
- maternal circulation, assessment of 400–401
- mature callus 230
- mechanical probes 6
- mechanical transducers 6
- meconium impaction 578f
- medial carpus 130–131, 131f, 133f
- medial collateral ligament (MCL) 64, 65f, 131, 195, 196f, 197f, 216f
 - hock 173
 - meniscal injuries 217f
- medial deep digital flexor tendon 180
- medial digital flexor tendon (MDFT) 91
- medial femoral condyle (MFC) 196f, 197f
 - caudal aspect of 204f
 - medial meniscus 204f
 - subchondral cyst-like lesions of 223f
- medial femoral epicondyle (MFE) 216f
- medial femorotibial joint 194–195, 197f, 224f
- medial lens subluxation 498f
- medial meniscus (MM) 204f, 217f, 218, 219f, 224f
 - chronic degenerative changes in 221f
 - cranial meniscotibial ligaments of 220f
- medial palmar intercarpal ligament (MPaICL) 134, 135f
- medial tibial condyle (MTC) 196f
- medial trochlear ridge (MTR) 192f, 210f, 211f
- medial window 91
- megaesophagus 534, 536f
- melanoma 506f, 514f, 515, 539f
- melting ulcer 491f
- meniscal injuries 215, 218, 220
- meniscofemoral ligament (mfl) 198f
- mesenteric abscess 563f
- mesenteric disruption 561
- mesothelioma 419f
- metacarpal condyle (MC) 59f
- metacarpophalangeal joint 61, 66, 73
- metacarpus (MC3) 51–52, 90
 - bony injuries 122–126

- extensor tendons 122
- palmar aspect 103*f*
- preparation and scanning technique 85–86
- scanning protocol for 86*t*
- tendinopathy/desmopathy. *See*
tendinopathy/desmopathy
- tendons and ligaments, appearance of 86–
91, 88*f*
- ultrasonographic anatomy 86–92, 87*f*
- metatarsus (MT3) 51–52. *See also* metacarpus
- scanning protocol for 86*t*
- metritis 567*f*
- micro-convex probes 7
- microconvex transducer 25, 27*f*, 39
- small footprint 25
- microphakia 486, 497*f*
- middle carpal joint (MCJ) 128
- Middle Cerebral Artery (MCA) 402–403
- middle patellar ligament (MiPL) 192, 192*f*,
193*f*, 195*f*, 204, 210, 211*f*
- distal third of 209*f*
- and trochlea 208*f*
- mineralization/ossification 77
- of meniscal fragment 218
- mirror-image artifact 17
- M-mode 4–5, 5*f*, 421, 425*f*, 429, 435*f*
- mucus/biofilm 339, 340*f*
- multicystic uterus masculinus 295*f*
- mural hematoma 563, 566*f*
- muscles 537
- atrophy 248
- brachiocephalic 537
- omohyoid 537
- sternocephalic 537
- sternohyoid and sternothyroid 537
- strap 537
- swellings in 537
- muscular capsule 285
- myocarditis 439*f*
- myositis, jugular vein 541*f*
- n**
- nasogastric tube 535*f*
- navicular bone 27*f*, 30
- flexor surface 40*f*
- navicular bursa 35*f*
- N-butylscopolammonium bromide 323
- neck:
- abscess 538*f*, 539*f*
- bones and joints 249–254
- of fetus 390, 390*f*
- jugular groove. *See* jugular groove
- lateral region 537, 539
- muscles 537, 538*f*
- muscles of 248
- preparation and scanning technique
510
- soft tissues 248–249, 248*f*
- ultrasound assessment 243
- upper cervical region. *See* upper cervical
region, neck
- negative frequency shift 17–18, 18*f*
- neonatal intensive care unit (NICU) 573
- neonate/neonatology:
- abdominal cavity 576–579
- intercostal spaces 573
- medical/physical condition 573
- patient 573
- rib fracture 412*f*
- size and lack of body fat 573
- thoracic cavity 574–576
- upper lung 573
- neoplasia 417
- neoplastic disease 420
- nephritis 455*f*
- perirenal edema and 458*f*
- nephrolith 455*f*, 456*f*, 457*f*
- nephrosplenic entrapment 474*f*
- non coronary cusp (NCC) 424*f*
- non-pregnant mare:
- in diestrus 351
- examination of 324
- reproductive tract 324
- ultrasonographic evaluation of 323
- uterine horn 394, 394*f*
- uterus 559
- non-septic tenosynovitis 74
- non-viable pregnancies 359*f*
- 33 to 39 days gestation 359*f*
- 24-day gestation 358*f*
- normal bladder 579*f*

O

oblique distal sesamoidean ligaments (ODSLs) 41–43, 45f

oblique transverse transsolar/cuneal ultrasound 34f

obturator artery 300

olecranon fossa 149

omohyoid muscle 537

omphalitis 578f

omphalophlebitis 578f

onion-ring lesion 100

operator error artifacts 11–12

Optical Coherence Tomography (OCT) 491f

optic nerve 484

orbital diameters 385–386

orchitis 313, 315f

osseous cyst-like lesions 61, 63f, 147f, 220

osseous lesions 161

osteoarthritis (OA) 29, 30f, 204

 of elbow joint 149, 151f

 femorotibial joint 220–221

 fetlock 59

 hip joint 237f

 of proximal interphalangeal joint 47

osteochondral fragmentation 61–64, 145f

osteochondritis dissecans (OCD) 60, 64f

osteochondrosis (OC) 59–61, 60, 61f, 62f, 183, 184f, 206, 209

 trochlear ridge 206

osteochondrosis dissecans (OCD) 206

osteomyelitis 184f

osteophytes 59, 204

ovarian hematomas 334–335, 335f

ovary/ovaries

 accessory corpora lutea on 356f

 anovulatory follicles 332–333, 332f

 contralateral 333, 334

 with corpus luteum 329f

 during early transitional phase 325f

 epithelial inclusion cysts 334, 334f

 granulosa cell tumor 333–334, 333f

 ovarian hematomas 334–335, 335f

 parovarian cysts 334, 334f

over-strain midsubstance injuries 77, 77f

ovulation date 357

P

PAL. *See* palmar/plantar annular ligament

palmar carpus 131–134, 134f, 135f

palmar foot 26f

palmar metacarpophalangeal ligament 65–66

palmar/plantar annular ligament (PAL) 44, 72, 72f, 75f, 76f

 digital flexor tendons sheath 76–77

palmar/plantar P1 fragments 67

palpation 323

 of internal reproduction tract 348f

per rectum 272, 558

 pregnancy diagnosis 347

papillary muscle ischemia 442f

paralumbar fossa (PLF) 446

paraphimosis 302, 307f

paratendinitis 106, 107f

parotidoauricularis muscle 256

parotid salivary gland 510–511, 513f, 514, 514f

parovarian cysts 334, 334f

parrot-beak tears 218

partial CL tears 215

pastern 39–47

 fetlock and 39

 preparation and scanning technique 41

 ultrasonographic abnormalities 42–47

 ultrasonographic anatomy 41–42, 42f

patellar fractures 209–210

 cranial aspect of 213f

patellar ligament (PL) 191, 194f, 210f

 injuries 204–206, 207f

 lateral 204

 middle 192, 192f, 193f, 195f, 204, 210, 211f

patellar luxation 225

patent urachus 555

patient preparation 10–11, 11f

peak systolic velocity (PSV) 302

pelvicitis 560, 560f

pelvic limb differences 91

pelvic urethra 273f–274f, 296f

 normal stallion 296

pelvis:

 areas of interest in 229

- dorsal sacroiliac ligaments 232, 234f, 235, 235f
- equipment 229
- hip joint 235–238
- ilium 229–230
- indications 229
- lumbar intertransverse joints 240, 241f
- lumbosacral joints 238–240, 239f, 240f
- per rectum* evaluation 238
- sacroiliac joint 240
- sacrum 240
- third trochanter 232, 233f, 234f
- tubera sacrale* 230, 230f, 231–232, 235f
- tuber coxa* 230, 230f, 231–232, 232f
- tuber ischium* 231–232, 232f
- ultrasound assessment 229
- penis:
 - anatomy 300
 - blood flow 302
 - body, proximal part of 303f
 - and cross-sections, diagram of 301f
 - distal part 301f, 306f, 308f
 - dorsal artery 300, 302f, 303f
 - free part 304f
 - glands 306f
 - in male 370
 - organizing hematoma of 309f
 - pathologies 302–309
 - ultrasound evaluation 300–302
 - urethra 302f
 - vascularization of 302
- percutaneous electrical nerve stimulation (PENS) 263
- periarticular injuries 67
- pericarditis 426
- periosteal new bone formation 144f
- periostitis 122
- periparturient hemorrhage 561
- peripheral pulses 388
- perirectal abscess 480f
- perirenal edema 458f
- perirenal hemorrhage 560, 560f
- peritoneal cavity 459–463
 - pathology of 560, 561
- peritoneal fluid 459, 559
- peritoneum 559
- peritonitis 459, 561–562, 562f, 563f, 564, 567f
 - bacterial 460f, 461f, 462f
 - exudative 462f
 - with fecal contamination 471f
 - fibrinous 461f
 - fibrin tags 462f
 - intestinal wall in 461f
 - jejunum in 461f
 - neonatal foal with 463f
 - transrectal view of 462f
- perivascular cellulitis/edema 544f
- perivascular injection 540
- peroneus tertius 177, 179f
 - rupture 177
 - tendon 225
- per rectum* 568
 - evaluation, pelvis 238
 - palpation 272, 558
- per vaginum* 568
- phased-array probes 7
- physics of ultrasound 1–5
 - B-mode 3–4
 - echogenicity 3–4
 - M-mode 4–5
 - ultrasound waves. *See* ultrasound waves, features of
- physis 183, 187f
 - distal tibial 185f
 - septic tibial 185f
- piezoelectric crystals 2
- placental dysfunction 399
- pleural abscess 415, 415f
- pleural effusion 413f, 414–415
- pleural fluid 560
- pleural hemorrhage 412
- pleural space 413–414, 414f
- pleuritis 413–414, 413f
 - fibrinous 413f
- pleuropneumonia 413f, 414f
- plugged ampullae 286
- pneumometa 568
- pneumonia 574f
- pneumopericardium 417
- pneumoperitoneum 461f, 564

- pneumothorax 13, 416–417, 416f
- pneumouterus 567f
- positive frequency shift 17, 18f
- post-anesthetic myopathy 155
- posterior chambers 483
 - fibrin accumulation in 485
- posterior lens luxation 499f
- post foaling complications
 - abdominal discomfort in 558
 - abdominal viscera, pathology of 563–568
 - bladder rupture 562
 - common variations 559
 - cystitis 562
 - diaphragmatic hernia 559–560, 560f
 - gastrointestinal tract 568–571
 - hemoperitoneum 561
 - normal anatomy 558–559
 - pelvicitis 560, 560f
 - peritoneal cavity, pathology of 560, 561
 - peritonitis 561–562, 562f, 563f
 - prepubic tendon 559
 - rectus abdominis 559
 - retroperitoneal hemorrhage (perirenal) 560
 - ruptured body wall 559
 - technique 558
 - transabdominal ultrasonography, limitations of 559
 - uroperitoneum 562
- post mating/breeding induced endometritis (PMIE/PBIE) 335
- post-partum lochia 564
- Potomac Horse Fever colitis 477f
- power Doppler 20
 - transverse trans-solar 41f
- pregnancy
 - abnormal conceptuses 359
 - database 383–384
 - day 21 366f
 - day 29 adjacent to cysts 365f
 - days 9 to 16 351
 - days 17 to 20 351, 353
 - days 21 to 28 353
 - days 30 to 40 353–354
 - days 40 to 60 354, 357–358
 - detectable heartbeat 353, 355f
 - early pregnancy examination 358
 - elective fetoplacental evaluation 384
 - embryonic stage 349
 - in equine practice 384
 - failure 358–360
 - gestational age of 357
 - loss and return 360f
 - loss prior to day 359
 - monitoring 383
 - non-viable 359f
 - resorbing 353f
- pregnancy diagnosis
 - detection 347
 - first 60 days 347
 - palpation 347
 - transrectal technique 347–349
- pregnant mare serum gonadotrophin (PMSG) 339, 356f
- preovulatory follicle 328f
- prepubic tendon 559
- preputial tissue 377
- priapism (persistent erection) 302
- primary tenosynovitis 73–76
- prostaglandin, release after twin
 - reduction 367
- proximal digital annular ligament (PDAL) 42, 46f
- proximal digital sheath 70, 70f, 75
- proximal interphalangeal joint (PIP) 42
 - abnormalities 47
 - articular cartilage of 47
 - lateral axial palmar ligament of 46f, 47f
 - osteoarthritis of 47
- proximal scutum (scutum proximale) 52, 72
 - abnormalities 82–84
- proximal sesamoid bones (PSBs) fractures 65, 66–67
- proximal suspensory desmitis 107–113, 109f, 110f, 111f
- pubis, fracture of 239f
- pulmonary abscesses 418
- pulmonary atelectasis 417, 417f
- pulmonary consolidation 416f, 417
- pulmonary edema 13
- pulmonary parenchyma 411f, 412, 574f

perpendicular 411f
 pulsed wave (PW) Doppler 18–19, 19f, 421, 429
 pus (ipopion) 485
 pyothorax 415, 415f

q

quadriceps muscle 191, 213f
 tears 211, 214f
 quantitative assessment, of flexor tendon and suspensory ligament size 91–92

r

radial tears 218
 radiography 43, 128, 214, 220
 raisin heart 428
 record-keeping systems 335
 rectal ampulla 384
 rectal trauma 560
 rectum 480
 rectus abdominis 559
 rectus femoris muscle 214f
 recurrence 101, 102f
 reflection 2–3
 refraction 3
 relative wall thickness (RWT) 428
 renal cortex 446
 renal dysplasia 456f
 retained placenta, 566, 568, 568f
 retina 484
 retinaculum of the carpus (RetC) 131, 132
 retina detachment 487, 502f, 503f
 retrobulbar cellulite 487
 retrobulbar diseases 487
 retrobulbar space 482
 retroperitoneal hemorrhage (perirenal) 560
 retropharyngeal lymph nodes 511, 524f
 abscess 523f, 524f
 reverberation artifacts 12–14, 12f, 13f, 488, 488f
Rhodococcus equi infection 249, 462f
 rib fractures 412, 412f, 413f
 right coronary cusp (RCC) 424f
 right dorsal colitis 475, 475f, 476f
 right dorsal colon (RDC) 446, 447f, 448f
 right hemithorax 410–411

right kidney 446, 558
 normal 454f
 right ovary 331
 right ventricular outflow tract (RVOT) 424f
 ring-down artifact 12, 13, 14f

s

sabulous cystitis 564f
 sacroiliac joint 229, 240
 caudal aspect of 241f
 sacrum 240
per rectum 242f
 salivary duct (Stensen's duct) 511
 salivary glands 262
 scapula 155–157
 fractures 155, 157, 159f
 scapulohumeral joint (SHJ) 167–172, 169f, 170f, 171f
 level 167f
 normal proximodistal ultrasound scans of 169f
 scar tissue 563
 scatter 3
 scintigraphy 57
 sclera 484
 scoring system 335
 scrotal content:
 echotexture of 313
 epididymis 313
 evaluation 311f
 qualitative evaluation of 313–314
 spermatic cord 313
 testis 313
 vaginal cavity 313, 314
 scrotum 310
 “seagull's wings” 487
 secretion 271
 sedation 148, 255
 seminal vesiculitis 289f, 290, 295
 sepsis 104
 septic arthritis 67–68, 149, 221, 225
 of elbow 150f
 iatrogenic 224f
 septic extensor carpi radialis tendon sheath 138f
 septic intraluminal fluid 566f

- septic pedal osteitis 36f
- septic synovitis 167, 177
- septic tendinitis 105f
- septic tenosynovitis 73
- serosanguinous peritoneal fluid 469f
- Sertoli cell tumor 316f
- sesamoid bones 90
- short lateral collateral ligament (SLCL) 174, 176f
- short medial collateral ligament (SMCL) 174f
- shoulder
 - normal point of 161f
 - scanning technique 155
 - scapula and supraglenoid tubercle 155–157
 - septic synovitis of 167
- side-lobe artifact 16–17, 17f
- slice-thickness artifact 15–16, 16f
- small intestinal muscular hypertrophy 472f
- small intestine 466–473, 469f
 - entrapped 571f
 - intussusception 577f
 - lipoma 470f, 471f
 - non-strangulating entrapment of 471f
 - strangulation 470f
 - volvulus 577f
 - volvulus with ventral sedimentation 470f
- soft tissues
 - neck 248–249, 248f
 - structures 255
 - swelling 175
- solar corium 33f
- SonoSite M-Turbo® ultrasound system 348
- spectral Doppler 18, 18f
- sperm accumulators 290
- spermatic cord 313
 - affected 315f
 - cross-section of 314f
 - hyperechoic inconsistencies in 315f
 - torsion 314f, 315f
- spleen 3, 4f, 410, 446, 558
 - cholangiohepatitis 450f, 451f
 - hemangiosarcoma 453f
 - lymphosarcoma 453f
 - normal 452f
 - splenic tear 453f
 - spleen–stomach relationship 452f, 453f, 466f
 - splenic vein 446
- spontaneous echo contrast (SEC) 437f
- spontaneous fetal movement 357
- squamous cell carcinoma (SCC) 507f
- stallion
 - internal genitalia 272, 286
 - internal reproductive tract 295
 - internal vaginal rings 272
 - pelvic urethra 272
 - penis. *See* penis
 - Tennessee Walking Horse 449f
 - testes size 319
- standoff pad 7, 8f, 49, 85, 148, 155, 482
- sterile ultrasound coupling gel 482
- sternocephalic muscle 537
- sternothyroid muscles 537
- stifle
 - abscesses 213f
 - anatomy of 191–203
 - femoropatellar joint. *See* femoropatellar joint
 - femorotibial joints. *See* femorotibial joints
 - lateral femorotibial joint 195, 198f, 199f
 - medial aspect 214f, 219f, 225f
 - medial femoral condyle 196f, 197f
 - medial femorotibial joint 194–195, 195f
 - preparation and scanning
 - technique 189–191
- stomach 465, 466, 571f
 - cranial aspect of 467f
 - dorsocaudal displacement 570
 - enlarged 576f
 - gastrointestinal tract 570
 - spleen and 466f
 - tube 467f
- straight distal sesamoidean ligaments (SDSLs) 41–43, 44f
 - insertion site of 43f
- strap muscles 537
- Streptococcus equi* 514, 523f
- Streptococcus equi zooepidemicus* 340f, 345f, 455f
- stress fractures 229, 243
 - in racehorses 231f
- stromal abscesses 484, 489f, 490f

- and anterior uveitis 492f
- stromal corneal lesion 493f
- Strongylus vulgaris* infection 540
- stylohyoid bone 514
- subacute synovitis 58f
- subchondral bone 206
 - defects 67
 - injury 57–59
- subchondral cyst-like lesions 220, 223f
- subcutaneous bursa 180
- subcutis 185
- subperiosteal hemorrhage 126f
- subsolar abscess 35f
- superficial digital flexor tendon (SDFT) 42,
 - 43, 69, 70, 72, 73, 76, 86, 89, 91, 96f, 100, 105f, 133f, 139f, 174, 175
 - complete rupture of 95
 - cross-sectional area of 97f
 - digital sheath abnormalities 80–82
 - and gastrocnemius tendon 181f
 - lateral branch of 80f
 - paratendinitis of 107f
 - proximal metacarpal 96f
 - tendinopathy/desmopathy 93–96, 96f
- superficial soft tissues 143
- supraglenoid tubercle 155–157
 - fracture 160f
- supraspinatus fossa 155, 159f, 166f
- supraspinatus muscle 163–166, 166f
- supraspinous ligament (SSpL) 243–245, 244f, 245f
- surgery 146
- suspensory desmopathy 106
- suspensory ligament (SL) 89–90, 106
 - branches 117–118, 118f, 119f
 - desmitis of body 113–117, 115f, 116f
 - size 91–92
- sustentaculum tali (ST) 182f, 183f, 187f
- swellings, in muscle 537
- synechiae 484
- synovial cavity 224f
- synovial effusion 203
- synovial fluid (SF) 177, 187, 254, 257f
- synovial fossa 149
- synovial hypertrophy 57

- synovial inflammation 214
- synovial membrane 61, 62f
- synovial plica 52
- synoviocoele 75, 75f
- synovitis 55–57, 177
- systolic left ventricular function 428–429

t

- tarsal sheath 177f, 180–181, 182f
- tarsocrural joint (TCJ) 174f
 - dorsal aspect 187f
 - dorsomedial pouch of 179f
- tarsometatarsal joint (TMTJ) 180
- tarsus
 - bony structures 181, 183
 - collateral ligaments 173–174
 - dorsal aspect 177, 186f
 - plantar aspect of 174–175
 - plantaromedial aspect 175–177
 - synovial structures 177–180
 - tarsal sheath 180–181
 - ultrasonography of 173
- tears 79–80
 - caudal horn 218
 - collateral ligament 215
 - cranial meniscotibial ligament 215
 - cranial meniscotibial ligaments (CMTL) 215
 - longitudinal 218
 - meniscal 215, 218, 220
 - splenic 453f
 - uterine 568f
 - vaginal 463f
- temporomandibular joint (TMJ) 255
 - abnormal 257f
 - caudolateral ultrasound image 256f
 - in normal adult horse 257f
 - normal anatomy and scanning technique 255–257
 - ultrasonographic abnormalities 257
- tendinitis/desmitis 92, 106
- tendinopathy/desmopathy 92–122
 - assessment of healing 97–100
 - chronic 100–102, 101f
 - cross-sectional area of 93

- deep digital flexor tendon 106
- general ultrasonographic changes 92–93
- local traumatic injuries 103–106, 103f
- semi-objective assessment of severity 96–97
- superficial digital flexor tendon 93–96, 96f
- tendons 86–91, 88f
 - biceps brachii 158f
 - carpus 143
 - of deep digital flexor tendon 177
 - hock 173–177
 - infraspinatus muscle and 166–167
 - and long digital extensor muscle 225
 - peroneus tertius 225
 - supraspinatus muscle and 163–166
 - triceps brachii 158f
- tensocopy 84
- testes. *See* testis
- testis
 - cranial–caudal view of 311f, 313f
 - height 311, 311f
 - interpretation 317, 319
 - lateral view 311f, 312f
 - length 311
 - measures 310–313
 - parenchyma 316f
 - probe type 310
 - scrotal content. *See* scrotal content
 - size 314
 - stallion position and location 310
 - tunica albuginea of 316f
 - varicocele 315f
 - volume, measurement and interpretation
 - of 314–319
 - width 311, 313, 313f
- third trochanter 232, 233f, 234f
- thoracic cavity (TC) 357f, 574–576
 - ventro-lateral aspect of 545f
- thoracic limb 88–89
- thoracic lymphadenopathy 414
- thoracic ultrasonography
 - diaphragm 410, 410f, 419f
 - diaphragmatic disruption 419
 - heart 409
 - hemothorax 414f, 415
 - increased pulmonary volume 419
 - interstitial pneumonia 419
 - left hemithorax 410
 - limitations of 420
 - musculature of 409
 - normal lung 410f
 - pleural effusion 413f, 414–415
 - pleural space, pathology of 413–414
 - pneumopericardium 417
 - pneumothorax 416–417
 - pulmonary abscesses 418, 418f
 - pulmonary appearance 411–412
 - pulmonary atelectasis 417
 - pulmonary consolidation 417
 - right hemithorax 410–411
 - technique 409
 - thoracic wall, pathology of 412–413
- thorax 13, 13f
- thrombophlebitis 540, 541f
 - anaerobic infection 543f
 - IV catheter 543f
 - of jugular vein 542f, 543f
- thrombosis, jugular 542f
- thyroid cartilage (TC) 259, 260f
- thyroid gland 520, 524f, 525
 - left lobe of 525f–527f
 - right lobe of 526f
- time-gain compensation (TGC) control 8, 9f, 15
- tissue Doppler imaging (TDI) 421, 429
- tissue harmonic imaging (THI) 9–10
- “T” ligament 25
- tongue 260–262
 - anatomy and scanning technique 260–261
 - longitudinal ultrasound image 261f
 - transverse ultrasound image 262f
 - ultrasonographic assessment 261
- total testicular volume (TTV) 317, 319f
- trabeculae:
 - of corpus cavernosum 304f, 308f, 309f
 - in corpus spongiosum 308f, 309f
- trace 18
- trachea 466f, 532, 534
 - with fluid 534f
 - normal 533f
- transabdominal ultrasonography 384

- limitations of 559
- transcorneal ultrasound 482
- transducer:
 - frequency 5–6, 6f
 - selection 10
 - types of 6–7
- trans-eyelid scan 483
- trans-palpebral ultrasound 482
- transrectal ultrasonography (TRUS)
 - 272, 384
- transsolar and transcuneal
 - ultrasonography 29–39
- transverse facial vein 257
- transverse ultrasound scan 29f
- traumatic arthritis 57–59
- triceps brachii muscle 155
 - on olecranon tuberosity 158f
- T-type linear probe 310
- tubera sacrale* 230, 230f, 231–232
 - asymmetry of 235f
- tuber coxa* 230, 230f, 231–232, 232f
 - fracture of 233f
- tuber ischium* 231–232, 232f
- tubular diverticula 271
- tunica albuginea 308
- turbulent flow 19
- twinning. *See also* twins
 - avoidance 363
 - ovulation induction products 363
- twins
 - flunixin meglumine 366
 - management 363
 - reduction by intracardiac/intra-thoracic
 - injection 384
 - reduction considerations 363
 - separating 364
 - success rate 368
 - 28-day adjacent 367f
- twin vesicles 364f
 - of different diameters 365f
 - differing sizes 364f
 - separated 364f
- two-dimensional (2-D) imaging 421
- 2-D speckle tracking (2DST) 421
- typhlitis 479f

u

- ulnaris lateralis muscle 150, 155, 156f, 157f
- ulnaris lateralis tendon (ULT) 130
- ultrasonography 1
- ultrasound 226
 - definition 2
 - guided procedures, head 263–264
 - physics 1–5
- ultrasound gel 310
- ultrasound machine 4
 - console 5
 - controls 7–10
 - depth 8
 - focus 8–9, 9f
 - frame rate 9
 - gain control 8
 - image optimization 7–10
 - image resolution 5–6, 6f
 - patient preparation 10–11, 11f
 - presets 7–8
 - time-gain compensation control 8, 9f
 - tissue harmonic imaging 9–10
 - transducer frequency 5–6, 6f
 - and transducer selection 10
 - transducer, types of 6–7
- ultrasound waves, features of 2–3
 - absorption 3
 - acoustic impedance 2–3, 2t
 - reflection 2–3
 - refraction 3
 - scatter 3
- umbilical arteries 550, 553f, 554f, 579f
- umbilical circulation 401
- umbilical cord (UC) 356f, 373, 376f, 396–399
 - of fetus 217-days-old 399f
 - and hippomane 397f
 - in male 370
 - at 180 days gestation 397f
 - placental root of 397f
 - 234 days gestation 398f
- umbilical hernia 555, 556f, 557, 557f
- umbilical remnant 548f, 550f
 - external 547, 549f
 - internal 555f
 - probe orientation 548f

- umbilical arteries 550, 553*f*
- umbilical hernia 555, 556*f*, 557, 557*f*
- umbilical vein 547, 550
- urachus 550, 555
- umbilical vein 547, 550, 551*f*, 552*f*
 - fetus 550
 - infection 550
 - linea alba 551*f*
 - omphalophlebitis 578*f*
 - pathology 550, 551*f*
- unidentified mass, in upper cervical region 527*f*–528*f*
- upper cervical region, neck 510–525
 - carotid artery 518
 - laryngeal lymph node 522*f*
 - left 512*f*
 - lymph nodes 520, 521*f*–522*f*
 - mandibular salivary gland 518
 - parotid salivary gland 510–511, 513*f*, 514, 514*f*
 - sites and associated structures in 512*f*
 - thyroid gland 520, 524*f*, 525, 525*f*–527*f*
 - ultrasound landmarks of 512*f*
 - unidentified mass 525, 527*f*–528*f*
- urachus 550, 555, 578*f*
 - infection 555
 - omphalitis 578*f*
 - patent 555
- urethra 300
 - corpus spongiosum of 302
 - penis 302*f*
- urethralis muscle 271
- urethral sinus 300, 307*f*
- urinary calculi 460*f*
- urine-filled bladder 562
- urogenital sinus 271
- uoperitoneum 460*f*, 562, 579*f*
- urovagina 343–345
- uterine arteries (UA) 400
 - rupture 339
- uterine body 331, 340*f*, 348
 - cross-section of 325*f*
 - diestrus 324*f*
 - estrus with edema 324*f*
 - gentle ballotment of 384
 - longitudinal ultrasonography 350*f*
 - lumen of 361*f*
- uterine cysts 350
 - cluster of 351*f*
- uterine edema 326, 568*f*
- uterine fluid, scoring system for 335
- uterine horn 348
 - cranial aspect 350*f*
 - in cross-section 341*f*
 - cross-section of 349*f*
 - 11-day embryonic vesicle in 351*f*
 - 15-day embryonic vesicle in 352*f*
 - 14-day embryonic vesicle in 352*f*
 - inverted 566, 566*f*
 - non-pregnant 394, 394*f*
 - 12-day embryonic vesicle in 351*f*
 - ultrasonography 350*f*
- uterine lumen 336*f*, 337*f*, 348*f*
- uteroplacental unit 394
- uterus 326, 335–342
 - air in 339, 339*f*
 - body of 324
 - broad ligament hematoma, 563, 565*f*
 - diestrus 329*f*, 335
 - with edema and normal fluid 326*f*
 - endometrial cups 339, 341*f*
 - endometrial cysts 335, 337*f*, 338*f*, 339
 - good visualization of 324*f*
 - horn 324*f*
 - intraluminal hemorrhage 567*f*
 - intrauterine fluid 335, 343*f*, 564
 - inverted uterine horn 566, 567*f*
 - involuting 566*f*
 - lumen of 324*f*, 325*f*
 - marbles 339, 342*f*
 - mucus/biofilm 339, 340*f*
 - mural hematoma 563, 566*f*
 - mural laceration, necrosis, and perforation 564
 - non-pregnant 559
 - pneumometra 568
 - post-partum 566
 - retained placenta 566, 568, 568*f*
 - uterine artery rupture 339
 - vaginal fluid 568

utriculus masculinus 271
 in colliculus masculinus 286
 uveitis 485, 490*f*, 492*f*, 496*f*
 cataracts and 497*f*

V

vagina 342–345
 and cervix 345*f*
 vault 345*f*
 vaginal cavity 313, 314
 fluid (hydrocele) in 318*f*
 vaginal fluid 569*f*
 uterus 568
 vaginal tear 463*f*
 valvular regurgitation 433–444
 varicocele 315*f*
 vasa deferentia 271, 273. *See also* vas deferens
 ampullae of 276*f*–280*f*, 294*f*, 295*f*
 vascular structures 540
 distal limb vasculature 540
 iliac arteries and aorta 540, 546
 jugular vein. *See* jugular vein
 lateral thoracic vein 540
 vas deferens. *See also* vasa deferentia
 ampulla of 283*f*–284*f*, 289
 dilated ampulla of 285*f*
 occluded ampulla of 285*f*
 terminal portion of 281*f*–282*f*

ventral cervix 344*f*
 ventral colon 559
 ventrolateral larynx 259
 vesicular glands 271
 ampulla of 281*f*
 enlarged 298
 fundus of 288*f*
 mucous membrane 271
 palpation 272
 secretion 280
 stallion with seminal vesiculitis 290*f*
 terminal portion of 281*f*
 ultrasound appearance 280
 ultrasound image of 289
 veterinary medicine 481
 veterinary ultrasonography 1
 villonodular synovitis 57
 vitreous chamber 484
 vitreous inflammation 486
 vitreous opacity 486–487

X

xiphoid cartilage 547
 xylazine 323

Y

yolk sac 350*f*
 envelopment of 357

